arXiv:1203.2677v4 [nlin.SI] 22 Jul 2012

On the τ-functions of the reduced Ostrovsky equation and the $A_2^{(2)}$ two-dimensional Toda system

Bao-Feng Feng[1], **Ken-ichi Maruno**[1] **and Yasuhiro Ohta**[2]

[1] Department of Mathematics, The University of Texas-Pan American, Edinburg, TX 78539-2999

[2] Department of Mathematics, Kobe University, Rokko, Kobe 657-8501, Japan

E-mail: `feng@utpa.edu`, `kmaruno@utpa.edu` and `ohta@math.kobe-u.ac.jp`

Abstract. The reciprocal link between the reduced Ostrovsky equation and the $A_2^{(2)}$ two-dimensional Toda system is used to construct the N-soliton solution of the reduced Ostrovsky equation. The N-soliton solution of the reduced Ostrovsky equation is presented in the form of pfaffian through a hodograph (reciprocal) transformation. The bilinear equations and the τ-function of the reduced Ostrovsky equation are obtained from the period 3-reduction of the B_∞ or C_∞ two-dimensional Toda system, *i.e.*, the $A_2^{(2)}$ two-dimensional Toda system. One of τ-functions of the $A_2^{(2)}$ two-dimensional Toda system becomes the square of a pfaffian which also become a solution of the reduced Ostrovsky equation. There is another bilinear equation which is a member of the 3-reduced extended BKP hierarchy. Using this bilinear equation, we can also construct the same pfaffian solution.

26 February 2013

PACS numbers: 02.30.Ik, 05.45.Yv

To be submitted to : *J. Phys. A: Math. Gen.*

1. Introduction

In this paper, we study the N-soliton solutions of the reduced Ostrovsky equation [1, 2]

$$\partial_x\left(\partial_t+u\partial_x\right)u-3u=0\,, \tag{1.1}$$

which is a special case ($\beta=0$) of the Ostrovsky equation

$$\partial_x\left(\partial_t+u\partial_x+\beta\partial_x^3\right)u-\gamma u=0\,. \tag{1.2}$$

The Ostrovsky equation was originally derived as a model for weakly nonlinear surface and internal waves in a rotating ocean [1, 2]. Later, the same equation was derived from different physical situations by several authors [2, 3, 4]. Note that the reduced Ostrovsky equation (1.1) is sometimes called the Vakhnenko equation or the Ostrovsky-Hunter equation [4, 5, 6, 7, 8]. Vakhnenko et al. constructed the N (loop) soliton solution of the reduced Ostrovsky equation by using a hodograph (reciprocal) transformation and the Hirota bilinear method [5, 6]. The same problem was approached from the point of view of inverse scattering method [7].

Differentiating the reduced Ostrovsky equation (1.1) with respect to x, we obtain

$$u_{txx}+3u_xu_{xx}+uu_{xxx}-3u_x=0\,, \tag{1.3}$$

which is known as the short wave limit of the Degasperis-Procesi (DP) equation [9, 10]. This equation is derived from the DP equation [11]

$$U_T+3\kappa^3U_X-U_{TXX}+4UU_X=3U_XU_{XX}+UU_{XXX}\,, \tag{1.4}$$

by taking a short wave limit $\varepsilon\to 0$ with $U=\varepsilon^2(u+\varepsilon u_1+\cdots)$, $T=\varepsilon^{-1}t$, $X=\varepsilon x$. Using this connection, Matsuno constructed the N-soliton solution of the short wave model of the DP equation, *i.e.*, the reduced Ostrovsky equation, from the N-soliton solution of the DP equation [10, 12, 13]. This N-soliton formula is equivalent to the one obtained by Vakhnenko. Hone and Wang pointed out that there is a reciprocal link between the reduced Ostrovsky equation and the first negative flow in the Sawada-Kotera hierarchy [9].

In this paper, we show the reciprocal link between the reduced Ostrovsky equation and the $A_2^{(2)}$ two-dimensional Toda system and investigate their τ-functions. Using this reciprocal link, we construct the N-soliton solution of the reduced Ostrovsky equation in the form of pfaffian. The bilinear equations and the τ-functions of the reduced Ostrovsky equation are systematically obtained from the period 3-reduction of the B_∞ or C_∞ two-dimensional Toda system. One of the τ-functions of the $A_2^{(2)}$ two-dimensional Toda system becomes the square of a pfaffian, by which N-soliton solution of the reduced Ostrovsky equation is expressed. We also show that another bilinear equation which is a member of the 3-reduced extended BKP hierarchy, or the so-called negative Sawada-Kotera hierarchy, can give rise to the reduced Ostrovsky equation by a hodograph transformation. Using this bilinear equation, we can also construct the same pfaffian solution.

2. The reduced Ostrovsky equation, the period 3-reduction of the B_∞ 2D-Toda system, and the 3-reduced extended BKP hierarchy

2.1. The two-dimensional Toda system of B_∞-type and its period 3-reduction

The two-dimensional Toda (2D-Toda) system of A_∞-type, which is also called the Toda field equation or the two-dimensional Toda lattice, is given as follows [14, 15, 16, 17]:

$$\frac{\partial^2\theta_n}{\partial x_1\partial x_{-1}}=-\sum_{m\in\mathbb{Z}}a_{n,m}e^{-\theta_m}\,,\quad n\in\mathbb{Z} \tag{2.1}$$

where the matrix $A=(a_{n,m})$ is the transpose of the Cartan matrix for the infinite dimensional Lie algebra A_∞ [18].

The A_∞ 2D-Toda system (2.1) may be written as

$$\frac{\partial^2\theta_n}{\partial x_1\partial x_{-1}}=e^{-\theta_{n-1}}-2e^{-\theta_n}+e^{-\theta_{n+1}}\,. \tag{2.2}$$

The A_∞ 2D-Toda system (2.1) is transformed into the bilinear equation

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}}-1\right)\tau_n\cdot\tau_n=\tau_{n-1}\tau_{n+1}\,, \tag{2.3}$$

through the dependent variable transformation

$$\theta_n = -\ln \frac{\tau_{n+1}\tau_{n-1}}{\tau_n^2}\,. \tag{2.4}$$

Here D_x is the Hirota D-operator which is defined as

$$D_x^n a(x)\cdot b(x) = (\partial_x - \partial_{x'})^n\, a(x)b(x')|_{x=x'}\,. \tag{2.5}$$

Lemma 2.1. *[Ueno-Takasaki[19], Babich-Matveev-Sall[20], Hirota[21],Nimmo-Willox[18]] The bilinear equations of the 2D-Toda lattice hierarchy including (2.3) have the following Gram-type determinant solution:*

$$\tau_n = \det\left(\psi_{i,j}^{(n)}\right)_{1\le i,j\le M}\,, \tag{2.6}$$

where

$$\psi_{i,j}^{(n)} = c_{i,j} + (-1)^n \int_{-\infty}^{x_1} \varphi_i^{(n)}\hat{\varphi}_j^{(-n)}\,dx_1\,. \tag{2.7}$$

Here $c_{i,j}$ are constants, $\varphi_{i,j}^{(n)}$ and $\hat{\varphi}_{i,j}^{(n)}$ satisfy

$$\frac{\partial \varphi_i^{(n)}}{\partial x_k} = \varphi_i^{(n+k)}\,, \quad \frac{\partial \hat{\varphi}_i^{(n)}}{\partial x_k} = (-1)^{k-1}\hat{\varphi}_i^{(n+k)}\,, \tag{2.8}$$

for $k = \pm 1, \pm 2, \pm 3, \cdots$.

For example, the linear independent set of functions $\{\varphi_i^{(n)}, \hat{\varphi}_j^{(n)}\}$ such that

$$\varphi_i^{(n)} = p_i^n e^{\xi_i}\,, \quad \hat{\varphi}_j^{(n)} = q_j^n e^{\eta_j}\,,$$

$\xi_i = p_i x_1 + \frac{1}{p_i}x_{-1} + p_i^2 x_2 + \frac{1}{p_i^2}x_{-2} + p_i^3 x_3 + \frac{1}{p_i^3}x_{-3} + \cdots + \xi_{i0}$ *and* $\eta_i = q_i x_1 + \frac{1}{q_i}x_{-1} - q_i^2 x_2 - \frac{1}{q_i^2}x_{-2} + q_i^3 x_3 + \frac{1}{q_i^3}x_{-3} + \cdots + \eta_{i0}$ *for $i, j = 1, 2, \cdots, M$, gives M-soliton solution of the A_∞ 2D-Toda system.*

Proof. See [21]. □

We impose the B_∞-reduction $\theta_n = \theta_{1-n}$ ($n \ge 1$) to the A_∞ 2D-Toda system (2.1), *i.e.*, fold the infinite sequence $\{\ldots, \theta_{-2}, \theta_{-1}, \theta_0, \theta_1, \theta_2, \ldots\}$ in the midpoint between θ_0 and θ_1 [22, 19, 18, 23]. Under this constraint, we have $\theta_0 = \theta_1$, $\theta_{-1} = \theta_2$, $\theta_{-2} = \theta_3$, $\ldots$.

For $n = 1$,

$$\begin{aligned}\frac{\partial^2\theta_1}{\partial x_1 \partial x_{-1}} &= e^{-\theta_0} - 2e^{-\theta_1} + e^{-\theta_2} \\ &= e^{-\theta_2} - e^{-\theta_1} \\ &= e^{-\theta_2} - 2e^{-(\theta_1+\ln 2)}\,.\end{aligned}$$

For $n = 2$,

$$\begin{aligned}\frac{\partial^2\theta_2}{\partial x_1 \partial x_{-1}} &= e^{-\theta_1} - 2e^{-\theta_2} + e^{-\theta_3} \\ &= 2e^{-(\theta_1+\ln 2)} - 2e^{-\theta_2} + e^{-\theta_3}\,.\end{aligned}$$

After redefining θ_1 by $\theta_1+\ln 2\to\theta_1$, we obtain the B_∞ 2D-Toda system [14, 17]:

$$\frac{\partial^2\theta_n}{\partial x_1\partial x_{-1}}=-\sum_{m\in\mathbb{Z}_{\geq 1}}a_{n,m}e^{-\theta_m}\,,\quad \text{for}\quad n\in\mathbb{Z}_{\geq 1}\,, \tag{2.9}$$

where the matrix $A=(a_{n,m})$ is the transpose of the Cartan matrix for the infinite dimensional Lie algebra B_∞ [18].

The B_∞ 2D-Toda system (2.9) is transformed into the bilinear equations

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}}-1\right)\tau_1\cdot\tau_1=\tau_1\tau_2\,, \tag{2.10}$$

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}}-1\right)\tau_n\cdot\tau_n=\tau_{n-1}\tau_{n+1}\,,\quad \text{for}\quad n\geq 2\,, \tag{2.11}$$

through the dependent variable transformation

$$\theta_1=-\ln\frac{\tau_2}{\tau_1}\,,\qquad \text{and}\qquad \theta_n=-\ln\frac{\tau_{n+1}\tau_{n-1}}{\tau_n^2}\quad \text{for } n\geq 2. \tag{2.12}$$

Lemma 2.2. *The bilinear equations of the B_∞ 2D-Toda system (2.10) and (2.11) have the N-soliton solution which is expressed as*

$$\tau_n=\det\left(\psi_{i,j}^{(n)}\right)_{1\leq i,j\leq 2N}\,, \tag{2.13}$$

where

$$\psi_{i,j}^{(n)}=c_{i,j}-2(-1)^n\int_{-\infty}^{x_1}\varphi_i^{(n)}\varphi_j^{(-n+1)}dx_1\,, \tag{2.14}$$

$$\varphi_i^{(n)}=p_i^n e^{\xi_i}\,,\quad \xi_i=p_ix_1+\frac{1}{p_i}x_{-1}+p_i^3x_3+\frac{1}{p_i^3}x_{-3}+\cdots+\xi_{i0}\,,$$

and $c_{i,j}=-c_{j,i}$, $c_{i,i}=0$.

Proof. Imposing the B_∞ reduction $\tau_n=\tau_{1-n}$, *i.e.*, folding the sequence of the τ-functions $\{\ldots,\tau_{-2},\tau_{-1},\tau_0,\tau_1,\tau_2,\tau_3,\ldots\}$ in the midpoint between τ_0 and τ_1, we have $\tau_0=\tau_1$, $\tau_{-1}=\tau_2$, $\tau_{-2}=\tau_3$, ... [22, 19, 18, 23]. Thus we obtain the bilinear equations (2.10) and (2.11) from the A_∞ 2D-Toda bilinear equation (2.3).

To impose the B_∞ reduction to the Gram-type determinant solution of the A_∞ 2D-Toda system, we impose the constraint $\hat{\varphi}_j^{(n)}=-2\frac{\partial}{\partial x_1}\varphi_j^{(n)}=-2\varphi_j^{(n+1)}$, $c_{i,j}=-c_{j,i}$, $c_{i,i}=0$, $M=2N$ and $x_{2k}\equiv 0$ for every interger k. With this constraint, each element of the Gram-type determinant has the following property:

$$\begin{aligned}\psi_{i,j}^{(n)}&=c_{i,j}-2(-1)^n\int_{-\infty}^{x_1}\varphi_i^{(n)}\varphi_j^{(-n+1)}dx_1=-c_{j,i}+2(-1)^{1-n}\int_{-\infty}^{x_1}\varphi_j^{(-n+1)}\varphi_i^{(n)}dx_1\\&=-\psi_{j,i}^{(1-n)}\,.\end{aligned} \tag{2.15}$$

Then the τ-function satisfies $\tau_n=\tau_{1-n}$. Therefore the N-soliton solution of the B_∞ 2D-Toda lattice is expressed by the above Gram-type determinant. □

Definition 2.3. *Let $A=(a_{i,j})_{1\leq i,j\leq 2N}$ be a $2N\times 2N$ skew-symmetric matrix. The pfaffian of A is defined by*

$$\mathrm{pf}(A)=\mathrm{pf}(a_{i,j})_{1\leq i,j\leq 2N}=\mathrm{pf}(1,2,...,2N)=\sum_{\substack{i_1<i_2<\cdots<i_N\\ i_1<j_1,...,i_N<j_N}}\mathrm{sgn}(\pi)a_{i_1,j_1}a_{i_2,j_2}\cdots a_{i_N,j_N}\,,$$

where $\pi = \begin{pmatrix} 1 & 2 & \cdots & 2N-1 & 2N \\ i_1 & j_1 & \cdots & i_N & j_N \end{pmatrix}$ *is a permutation of* $\{1,2,...,2N-1,2N\}$.

The pfaffian can be computed recursively by

$$\mathrm{pf}(1,2,...,2N-1,2N) = \sum_{i=2}^{2N} (-1)^i \mathrm{pf}(1,i)\mathrm{pf}(2,3,...,i-1,i+1,...,2N-1,2N)\,.$$

For a skew-symmetric matrix A, we have $[\mathrm{pf}(A)]^2 = \det(A)$.

Lemma 2.4. *The τ-function* τ_1 *of the bilinear equations (2.10) and (2.11) is written in the form of pfaffian*

$$\tau_1 = \tau^2\,, \qquad \tau = \mathrm{pf}(1,2,...,2N-1,2N)\,, \tag{2.16}$$

where $\mathrm{pf}(i,j) = c_{i,j} + \int_{-\infty}^{x_1} D_{x_1}\varphi_i^{(0)} \cdot \varphi_j^{(0)} dx_1$, $\varphi_i^{(n)} = p_i^n e^{\xi_i}$, $\xi_i = p_i^{-1}x_{-1} + p_i x_1 + p_i^3 x_3 + \frac{1}{p_i^3}x_{-3} + \cdots + \xi_i^0$, $c_{i,j} = -c_{j,i}$, $c_{i,i} = 0$.

Proof. See [21, 24]. □

Lemma 2.5. *The bilinear equations*

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}} - 1\right)\tau_1 \cdot \tau_1 = \tau_1\tau_2\,, \tag{2.17}$$

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}} - 1\right)\tau_2 \cdot \tau_2 = \tau_1^2\,, \tag{2.18}$$

have the N-soliton solution which is expressed as

$$\tau_n = \det\left(\psi_{i,j}^{(n)}\right)_{1\le i,j\le 2N}\,, \tag{2.19}$$

where

$$\psi_{i,j}^{(n)} = c_{i,j} + \frac{2p_i}{p_i + p_j}\left(-\frac{p_i}{p_j}\right)^{n-1} e^{\xi_i + \xi_j}\,, \qquad \xi_i = p_i x_1 + \frac{1}{p_i}x_{-1} + \xi_{i0}\,,$$

and $c_{i,j} = \delta_{j,2N+1-i}c_i$, $c_i = -c_{2N+1-i}$, $p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 0$.

Proof. Imposing a period 3-reduction $\tau_n = \tau_{n+3}$ to the sequence of the τ-functions $\{\ldots,\tau_3,\tau_2,\tau_1,\tau_1,\tau_2,\tau_3,\ldots\}$, it becomes $\{\ldots,\tau_1,\tau_2,\tau_1,\tau_1,\tau_2,\tau_1,\ldots\}$ [22, 19, 18, 23]. Thus we obtain two bilinear equations (2.17) and (2.18).

In Lie algebraic terms, this corresponds to the reduction to the affine Lie algebra $A_2^{(2)}$ from the infinite dimensional Lie algebra B_∞ [22, 19, 18, 23].

To impose a period 3-reduction to the N-soliton solution, we add a constraint $p_i^3 + p_{2N+1-i}^3 = 0$ ($p_i \neq -p_{2N+1-i}$) for $i = 1,2,\cdots,N$ and $c_{i,j} = \delta_{j,2N+1-i}c_i$, $c_i = -c_{2N+1-i}$. To show $\tau_{n+3} = \tau_n$, we manipulate τ_n as follows:

$$\begin{aligned}
\tau_n &= \det\left(\delta_{j,2N+1-i}c_i + \frac{2p_i}{p_i+p_j}\left(-\frac{p_i}{p_j}\right)^{n-1} e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= e^{2\sum_{k=1}^{2N}\xi_k} \det\left(\delta_{j,2N+1-i}c_i\left(-\frac{p_j}{p_i}\right)^{n-1} e^{-\xi_i-\xi_j} + \frac{2p_i}{p_i+p_j}\right)_{1\le i,j\le 2N} \\
&= e^{2\sum_{k=1}^{2N}\xi_k} \det\left(\delta_{j,2N+1-i}c_i\left(-\frac{p_{2N+1-i}}{p_i}\right)^{n-1} e^{-\xi_i-\xi_{2N+1-i}} + \frac{2p_i}{p_i+p_j}\right)_{1\le i,j\le 2N}\,.
\end{aligned}$$

Since $\left(-\frac{p_{2N+1-i}}{p_i}\right)^3 = 1$, the relation $\tau_{n+3} = \tau_n$ is satisfied. □

Letting $f = \tau_1$ and $g = \tau_2$, we obtain the bilinear equations

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}} - 1\right) f \cdot f = fg\,, \tag{2.20}$$

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}} - 1\right) g \cdot g = f^2\,. \tag{2.21}$$

Corollary 2.6. *The τ-function $f = \tau_1$ of the bilinear equations (2.20) and (2.21) is written in the form of pfaffian*

$$f = \tau_1 = \tau^2\,, \qquad \tau = \mathrm{pf}(1,2,\ldots,2N-1,2N)\,, \tag{2.22}$$

where

$$\mathrm{pf}(i,j) = c_{i,j} + \frac{p_i - p_j}{p_i + p_j}e^{\xi_i+\xi_j}\,, \qquad \xi_i = p_i x_1 + \frac{1}{p_i}x_{-1} + \xi_{i0}\,,$$

and $c_{i,j} = \delta_{j,2N+1-i}c_i$, $c_i = -c_{2N+1-i}$, $p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 0$.

Proof. Using Lemma 2.4 and 2.5, we obtain the above formula. □

Let $k_i = p_i + p_{2N+1-i}$. From $p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 0$, we obtain $p_i = \frac{1}{6}(3k_i + \mathrm{i}\sqrt{3}k_i)$, $p_{2N+1-i} = \frac{1}{6}(3k_i - \mathrm{i}\sqrt{3}k_i)$, $p_i p_{2N+1-i} = \frac{k_i^2}{3}$ and $\frac{1}{pi} + \frac{1}{p_{2N+1-i}} = \frac{3}{k_i}$. Thus

$$\xi_i + \xi_{2N+1-i} = k_i x_1 + \frac{3}{k_i}x_{-1} + \xi_{i0} + \xi_{2N+1-i0}\,.$$

In the pfaffian solution, all phase functions can be expressed by the summation of $\xi_i + \xi_{2N+1-i}$. So the phase functions can be expressed by the parameters $\{k_i\}$ $(i = 1,2,\cdots,N)$. Each coefficient of exponential functions can be normalized to 1 after absorption into phase constants or can be rewritten by the parameters $\{k_i\}$. Thus it is possible to rewrite the above τ-function by using the parameters $\{k_i\}$ instead of $\{p_i\}$.

Examples:

For $N = 1$,

$$\tau = \mathrm{pf}(1,2) = c_1 + \frac{p_1 - p_2}{p_1 + p_2}e^{\xi_1+\xi_2}\,. \tag{2.23}$$

Letting $c_1 = 1$ and $e^{\gamma_1} = \frac{p_1-p_2}{p_1+p_2}$, the τ-function can be rewritten as

$$\tau = \mathrm{pf}(1,2) = 1 + e^{\xi_1+\xi_2+\gamma_1}\,, \tag{2.24}$$

where

$$\xi_1 + \xi_2 = k_1 x_1 + \frac{3}{k_1}x_{-1} + \xi_{10} + \xi_{20}\,.$$

For $N = 2$,

$$\begin{aligned}\tau &= \mathrm{pf}(1,2,3,4) = \mathrm{pf}(1,2)\mathrm{pf}(3,4) - \mathrm{pf}(1,3)\mathrm{pf}(2,4) + \mathrm{pf}(1,4)\mathrm{pf}(2,3)\\ &= \frac{p_1 - p_2}{p_1 + p_2}e^{\xi_1+\xi_2}\frac{p_3 - p_4}{p_3 + p_4}e^{\xi_3+\xi_4} - \frac{p_1 - p_3}{p_1 + p_3}e^{\xi_1+\xi_3}\frac{p_2 - p_4}{p_2 + p_4}e^{\xi_2+\xi_4}\end{aligned}$$

$$
\begin{aligned}
&\quad + \left(c_1 + \frac{p_1-p_4}{p_1+p_4} e^{\xi_1+\xi_4}\right)\left(c_2 + \frac{p_2-p_3}{p_2+p_3} e^{\xi_2+\xi_3}\right) \\
&= c_1c_2 + c_2\frac{p_1-p_4}{p_1+p_4} e^{\xi_1+\xi_4} + c_1\frac{p_2-p_3}{p_2+p_3} e^{\xi_2+\xi_3} \\
&\quad + \left(\frac{p_1-p_2}{p_1+p_2}\frac{p_3-p_4}{p_3+p_4} - \frac{p_1-p_3}{p_1+p_3}\frac{p_2-p_4}{p_2+p_4} + \frac{p_1-p_4}{p_1+p_4}\frac{p_2-p_3}{p_2+p_3}\right) e^{\xi_1+\xi_2+\xi_3+\xi_4}\,.
\end{aligned}
$$

Letting $c_1 = c_2 = 1$ and $e^{\gamma_1} = \frac{p_1-p_4}{p_1+p_4}$ and $e^{\gamma_2} = \frac{p_2-p_3}{p_2+p_3}$, the above τ-function becomes

$$
\tau = 1 + e^{\xi_1+\xi_4+\gamma_1} + e^{\xi_2+\xi_3+\gamma_2} + b_{12} e^{\xi_1+\xi_2+\xi_3+\xi_4+\gamma_1+\gamma_2}\,,
$$

where

$$
\begin{aligned}
b_{12} &= \frac{p_1-p_2}{p_1+p_2}\frac{p_3-p_4}{p_3+p_4}\frac{p_1+p_4}{p_1-p_4}\frac{p_2+p_3}{p_2-p_3} - \frac{p_1-p_3}{p_1+p_3}\frac{p_2-p_4}{p_2+p_4}\frac{p_1+p_4}{p_1-p_4}\frac{p_2+p_3}{p_2-p_3} + 1 \\
&= \frac{(k_1-k_2)^2(k_1^2-k_1k_2+k_2^2)}{(k_1+k_2)^2(k_1^2+k_1k_2+k_2^2)}\,,
\end{aligned}
$$

$$
\xi_1+\xi_4 = k_1x_1 + \frac{3}{k_1}x_{-1} + \xi_{10} + \xi_{40}\,, \quad \xi_2+\xi_3 = k_2x_1 + \frac{3}{k_2}x_{-1} + \xi_{20} + \xi_{30}\,.
$$

The N-soliton solution is written in the following form:

$$
\tau = \sum_{\mu=0,1} \exp\left[\sum_{i=1}^{N} \mu_i\eta_i + \sum_{i<j}^{(N)} \mu_i\mu_j \ln b_{ij}\right], \tag{2.25}
$$

$$
b_{ij} = \frac{(k_i-k_j)^2(k_i^2-k_ik_j+k_j^2)}{(k_i+k_j)^2(k_i^2+k_ik_j+k_j^2)} \quad \text{for} \quad i<j\,, \quad \eta_i = \xi_i + \xi_{2N+1-i} = k_ix_1 + \frac{3}{k_i}x_{-1} + \eta_{i0}\,,
$$

where $\sum_{\mu=0,1}$ means the summation over all possible combinations of $\mu_i = 0$ or 1 for $i = 1,2,\cdots,N$, and $\sum_{i<j}^{(N)}$ means the summation over all possible combinations of N elements under the condition $i<j$. This is consistent with the results in [5, 6].

Lemma 2.7. *The τ-function f of (2.20) and (2.21) gives the solution of the reduced Ostrovsky equation through the dependent variable transformation*

$$
u = -(\ln f)_{x_1x_1} = -2(\ln\tau)_{x_1x_1}\,,
$$

and the hodograph (reciprocal) transformation

$$
\left\{
\begin{aligned}
x &= x_{-1} + \textstyle\int_{-\infty}^{x_1} u(x_1', x_{-1})\,dx_1' \\
&= x_{-1} - (\ln f)_{x_1}\,, \\
t &= x_1\,.
\end{aligned}
\right. \tag{2.26}
$$

Proof. The bilinear equation (2.20) is rewritten as

$$
-(\ln f)_{x_1x_{-1}} = \frac{g}{f} - 1\,. \tag{2.27}
$$

Let

$$
\rho = \frac{f}{g} = \frac{1}{1-(\ln f)_{x_1x_{-1}}}\,, \quad u = -(\ln f)_{x_1x_1}\,. \tag{2.28}
$$

Then (2.27) becomes

$$u_{x_{-1}} = \left(\frac{1}{\rho}\right)_{x_1} . \tag{2.29}$$

This is rewritten as

$$(\ln\rho)_{x_1} = -\rho u_{x_{-1}} . \tag{2.30}$$

The bilinear equations (2.20) and (2.21) are written as

$$-(\ln f)_{x_1 x_{-1}} + 1 = \frac{1}{\rho} , \tag{2.31}$$

$$-(\ln g)_{x_1 x_{-1}} + 1 = \rho^2 . \tag{2.32}$$

Subtracting (2.32) from (2.31), we obtain

$$-(\ln\rho)_{x_1 x_{-1}} = \frac{1}{\rho} - \rho^2 , \tag{2.33}$$

which leads to

$$\rho^3 = 1 + \rho(\ln\rho)_{x_1 x_{-1}} . \tag{2.34}$$

Using (2.30), it becomes

$$\rho^3 = 1 - \rho(\rho u_{x_{-1}})_{x_{-1}} . \tag{2.35}$$

Let us consider the hodograph (reciprocal) transformation

$$\left\{ \begin{array}{l} x = x_{-1} + \int_{-\infty}^{x_1} u(x_1', x_{-1}) dx_1' \\ \quad = x_{-1} - (\ln f)_{x_1} , \\ t = x_1 . \end{array} \right. \tag{2.36}$$

This yield

$$\left\{ \begin{array}{l} \frac{\partial x}{\partial x_{-1}} = 1 - (\ln f)_{x_1 x_{-1}} , \\ \frac{\partial x}{\partial x_1} = -(\ln f)_{x_1 x_1} , \end{array} \right. \tag{2.37}$$

and

$$\left\{ \begin{array}{l} \partial_{x_{-1}} = \frac{1}{\rho}\partial_x , \\ \partial_{x_1} = \partial_t + u\partial_x . \end{array} \right. \tag{2.38}$$

Applying the hodograph (reciprocal) transformation to (2.30) and (2.35), we obtain

$$\left\{ \begin{array}{l} (\partial_t + u\partial_x)\ln\rho = -u_x , \\ \rho^3 = 1 - u_{xx} . \end{array} \right. \tag{2.39}$$

This is equivalent to

$$(\partial_t + u\partial_x)\ln(1 - u_{xx}) = -3u_x , \tag{2.40}$$

which can be written as

$$(\partial_t + u\partial_x)(1 - u_{xx}) = -3u_x(1 - u_{xx}) . \tag{2.41}$$

This is nothing but the short wave limit of the DP equation which is equivalent to the reduced Ostrovsky equation. □

Remark 2.8. Setting $\rho = \frac{1}{\mathcal{H}}$ in (2.33), we obtain

$$(\ln \mathcal{H})_{x_1 x_{-1}} = \mathcal{H} - \frac{1}{\mathcal{H}^2}\,. \tag{2.42}$$

This is the Tzitzeica equation which is one of prime integrable systems that appears in classical differential geometry [25, 26, 23, 24]. This equation is sometimes called the Dodd-Bullough-Mikhailov equation since this was found independently by Dodd-Bullough and Mikhailov [27, 15, 16].

Theorem 2.9. *The N-soliton solution of the reduced Ostrovsky equation is given by the following formula:*

$$\begin{aligned} &u = -(\ln f)_{x_1 x_1} = -2(\ln \tau)_{x_1 x_1}\,,\\ &f = \tau^2\,, \qquad \tau = \mathrm{pf}(1, 2, ..., 2N-1, 2N)\,, \end{aligned} \tag{2.43}$$

where

$$\mathrm{pf}(i, j) = c_{i,j} + \frac{p_i - p_j}{p_i + p_j} e^{\xi_i + \xi_j}\,, \qquad \xi_i = p_i x_1 + \frac{1}{p_i} x_{-1} + \xi_{i0}\,,$$

and $c_{i,j} = \delta_{j,2N+1-i} c_i\,,\ c_i = -c_{2N+1-i}\,,\ p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 0\,,$

$$\left\{ \begin{array}{l} x = x_{-1} + \int_{-\infty}^{x_1} u(x_1', x_{-1}) dx_1' \\ \quad = x_{-1} - (\ln f)_{x_1}\,, \\ t = x_1\,. \end{array} \right. \tag{2.44}$$

Proof. From Corollary 2.6 and Lemma 2.7, we obtain this theorem. □

Corollary 2.10. *The N-soliton solution of the reduced Ostrovsky equation is given by the following formula:*

$$u = -2(\ln \tau)_{x_1 x_1}\,,$$

$$\tau = \sum_{\mu=0,1} \exp\left[\sum_{i=1}^{N} \mu_i \eta_i + \sum_{i<j}^{(N)} \mu_i \mu_j \ln b_{ij} \right]\,, \tag{2.45}$$

$$b_{ij} = \frac{(k_i - k_j)^2 (k_i^2 + k_i k_j + k_j^2)}{(k_i + k_j)^2 (k_i^2 - k_i k_j + k_j^2)} \quad \text{for} \quad i < j\,, \quad \eta_i = k_i x_1 + \frac{3}{k_i} x_{-1} + \eta_{i0}\,,$$

and

$$\left\{ \begin{array}{l} x = x_{-1} + \int_{-\infty}^{x_1} u(x_1', x_{-1}) dx_1' \\ \quad = x_{-1} - (\ln f)_{x_1}\,, \\ t = x_1\,. \end{array} \right. \tag{2.46}$$

Here $\sum\limits_{\mu=0,1}$ *means the summation over all possible combinations of* $\mu_i = 0$ *or* 1 *for* $i = 1, 2, \cdots, N$, *and* $\sum\limits_{i<j}^{(N)}$ *means the summation over all possible combinations of N elements under the condition* $i < j$.

Note that the form of the N-soliton solution in Corollary 2.10 is equivalent to the one obtained by Morrison, Parkes and Vakhnenko in [6].

Remark 2.11. It is well known that the 3-reduced B-Toda system is exactly the same as the 3-reduced C-Toda system. Thus the above τ-function can be obtained by the period 3-reduction of C_∞ 2D-Toda system. For more details, see Appendix.

2.2. The reduced Ostrovsky equation and the 3-reduced extended BKP hierarchy

Consider a member of the extended BKP hierarchy (the BKP hierarchy with negative time variables) [21, 22, 28, 29, 30, 31, 32]

$$[(D_{x_3}-D_{x_1}^3)D_{x_{-1}}+3D_{x_1}^2]\tau\cdot\tau=0\,, \tag{2.47}$$

which is the bilinear equation of a nonlinear partial differential equation

$$w_{x_{-1}x_3}-w_{x_1x_1x_1x_{-1}}-3(w_{x_1}w_{x_{-1}})_{x_1}+3w_{x_1x_1}=0\,, \tag{2.48}$$

through the dependent variable transformation $w=2(\ln\tau)_{x_1}$ [21].

The N-soliton solution of (2.47) is expressed in the form of pfaffian.

Lemma 2.12. *[Date-Jimbo-Kashiwara-Miwa [28], Hirota [21, 31, 32]] The bilinear equation (2.47) has the pfaffian solution*

$$\tau=\mathrm{pf}(1,2,...,2N-1,2N)\,,$$

where $\mathrm{pf}(i,j)=c_{i,j}+\int_{-\infty}^{x_1}D_{x_1}h_i(\mathbf{x})\cdot h_j(\mathbf{x})dx_1$, $c_{i,j}=-c_{j,i}$, $c_{i,i}=0$, $\mathbf{x}=(\ldots,x_{-3},x_{-1},x_1,x_3,x_5\ldots)$ *and* $h_i(\mathbf{x})$ *satisfies the linear differential equation*

$$\frac{\partial}{\partial x_n}h_i(\mathbf{x})=\frac{\partial^n}{\partial x_1^n}h_i(\mathbf{x})\,. \tag{2.49}$$

Here we define

$$\frac{\partial}{\partial x_{-1}}h_i(\mathbf{x})\equiv\int_{-\infty}^{x_1}h_i(\mathbf{x})dx_1\,, \tag{2.50}$$

for $n=-1$. *Particularly, we could choose*

$$h_i(\mathbf{x})=\exp(p_i^{-1}x_{-1}+p_ix_1+p_i^3x_3+\xi_i^0)\,. \tag{2.51}$$

Alternatively, the N-soliton solution of (2.47) can be expressed as

$$\tau=\sum_{\mu_i=0,1}\exp\left[\sum_{i=1}^{N}\mu_i\eta_i+\sum_{1\le i<j\le N}B_{i,j}\mu_i\mu_j\right]\,, \tag{2.52}$$

where $\sum_{\mu_i=0,1}$ denotes the summation over all possible combinations of $\mu_i=0,1$ for $i=1,2,\cdots,N$, and $\sum_{1\le i<j\le N}$ is the sum over all pairs i,j $(i<j)$ chosen from $\{1,2,\cdots,2N\}$. Here $\exp(\eta_i)$ is defined as

$$\exp(\eta_i)=\exp(\xi_i+\hat{\xi}_i)\,,$$

where $\xi_i = p_i^{-1}x_{-1} + p_i x_1 + p_i^3 x_3 + \xi_i^0$ and $\hat{\xi}_i = q_i^{-1}x_{-1} + q_i x_1 + q_i^3 x_3 + \hat{\xi}_i^0$, and $B_{i,j}$ is given by

$$e^{B_{i,j}} = b_{i,j} = \frac{(p_i - p_j)(p_i - q_j)(q_i - p_j)(q_i - q_j)}{(p_i + p_j)(p_i + q_j)(q_i + p_j)(q_i + q_j)}\,.$$

Proof. See [21]. ∎

Lemma 2.13. *The bilinear equation*

$$(D_{x_{-1}} D_{x_1}^3 - 3D_{x_1}^2)\tau \cdot \tau = 0\,, \tag{2.53}$$

has the the following N-soliton solution:

$$\tau = \mathrm{pf}(1, 2, \ldots, 2N-1, 2N)\,,$$

where

$$\mathrm{pf}(i,j) = c_{i,j} + \frac{p_i - p_j}{p_i + p_j} e^{\xi_i + \xi_j}\,, \qquad \xi_i = p_i x_1 + p_i^{-1} x_{-1} + \xi_{i0}\,,$$

and $c_{i,j} = \delta_{j,2N+1-i} c_i\,,\ c_i = -c_{2N+1-i}\,,\ p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 0$.

Proof. Since all phase functions are expressed by the summation of $\xi_i + \xi_{2N+1-i}$, x_3 is eliminated from the τ-function by imposing constraints: $p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 0$ for $i = 1, 2, \cdots, N$ and $c_{i,j} = \delta_{j,2N+1-i} c_i$, $c_i = -c_{2N+1-i}$ to the N-soliton solution of the BKP τ-function in Lemma 2.12 [22, 28, 29, 30]. Thus $\partial_{x_3}\tau = 0$ is satisfied. ∎

Lemma 2.14. *The reduced Ostrovsky equation (1.1) is transformed into the bilinear equation*

$$(D_{x_{-1}} D_{x_1}^3 - 3D_{x_1}^2)\tau \cdot \tau = 0\,, \tag{2.54}$$

through the dependent transformation

$$u = -w_{x_1} = -2(\ln \tau)_{x_1 x_1}\,, \tag{2.55}$$

and the hodograph (reciprocal) transformation (2.36).

Proof. Applying (2.38) to the reduced Ostrovsky equation (1.1), we obtain

$$\rho u_{x_1 x_{-1}} = 3u\,, \tag{2.56}$$

which yields

$$u_{x_1 x_{-1}} = \frac{3}{\rho} u\,. \tag{2.57}$$

Let us introduce w such that $u = -w_{x_1} = -(\ln f)_{x_1 x_1}$. From (2.31), we obtain

$$\frac{1}{\rho} = 1 - w_{x_{-1}}\,. \tag{2.58}$$

Using this relation, (2.57) can be rewritten as

$$w_{x_1 x_1 x_{-1}} + 3w_{x_{-1}} w_{x_1} - 3w_{x_1} = 0\,. \tag{2.59}$$

This equation is transformed into a bilinear equation

$$(D_{x_{-1}} D_{x_1}^3 - 3D_{x_1}^2)\tau \cdot \tau = 0\,, \tag{2.60}$$

through the dependent transformation

$$w = 2(\ln\tau)_{x_1}\,. \tag{2.61}$$

Note that the hodograph (reciprocal) transformation (2.36) can be expressed as

$$\begin{cases} x = x_{-1} - 2(\ln\tau)_{x_1}\,, \\ t = x_1\,. \end{cases} \tag{2.62}$$

□

From Lemma 2.13 and 2.14, we can also prove Theorem 2.9.

Remark 2.15. Vakhnenko and Parkes found the hodograph (reciprocal) transformation (2.62) and the bilinear equation (2.60). It is noted that we can switch x_{-1} and x_1, i.e., the alternative hodograph (reciprocal) transformation $x = x_1 - (\ln f)_{x_{-1}} = x_1 - 2(\ln\tau)_{x_{-1}}$, $t = x_{-1}$ can be used instead of the above hodograph (reciprocal) transformation.

Remark 2.16. Morrison, Parkes and Vakhnenko pointed out that the bilinear equation (2.60) is a special case of the Ito equation [6, 33]. Hone and Wang pointed out that the reduced Ostrovsky equation is linked into the first negative flow of the Sawada-Kotera hierarchy. The bilinear equation (2.60) is nothing but the bilinear equation of the first negative flow of the Sawada-Kotera hierarchy which can be obtained from the BKP hierarchy with the 3-reduction, *i.e.*, this is related to the $A_2^{(2)}$ affine Lie algebra [22].

Remark 2.17. The relationship between the Tzitzeica equation and the 3-reduced BKP hierarchy was pointed out by Lambert et al. [34].

3. Conclusions

In this paper, we have shown the reciprocal link between the reduced Ostrovsky equation and the $A_2^{(2)}$ 2D-Toda system. Using this reciprocal link, we have constructed the N-soliton solution of the reduced Ostrovsky equation in the form of pfaffian. The bilinear equations and the τ-function of the reduced Ostrovsky equation have been obtained from the period 3-reduction of the B_∞ or C_∞ 2D-Toda system. One of the τ-functions of the $A_2^{(2)}$ 2D-Toda system has been shown to be the square of a pfaffian, by which the N-soliton solution of the reduced Ostrovsky equation is represented. We have shown that the bilinear equation for a member of the 3-reduced extended BKP hierarchy (the first negative flow of the Sawada-Kotera hierarchy) can also give rise to the the reduced Ostrovsky equation, thus, the same pfaffian solution.

It should be noted that a similar approach in this paper was used to the short wave model of the Camassa-Holm equation in our previous paper [35]. We proposed an integrable semi-discrete and an integrable fully discrete analogues of the short wave model of the Camassa-Holm equation based on bilinear equations and a determinant solution. Thus we can construct integrable discrete analogues of the reduced Ostrovsky equation (or the short wave model of the DP equation) by using the result in this paper. We also would like to comment on the N-soliton solution of the DP equation can be obtained by the reduction of the 2D-Toda system developed in this paper. Our approach gives the N-soliton solution of the DP equation in a much simpler way compared to Matsuno's method [12, 13]. We will report the detail in our forthcoming paper.

Appendix: The period 3-reduction of the C_∞ 2D-Toda system

In this appendix, we show that the τ-function obtained in the section 2 can be obtained from the period 3-reduction of the C_∞ 2D-Toda system.

We impose the C_∞-reduction $\theta_n = \theta_{-n}$ $(n \geq 0)$ to the A_∞ 2D-Toda system, *i.e.*, fold the infinite sequence $\{\dots,\theta_{-2},\theta_{-1},\theta_0,\theta_1,\theta_2,\dots\}$ in θ_0 [22, 19, 18, 23]. Then we obtain the C_∞ 2D-Toda system [14, 17]:

$$\frac{\partial^2\theta_n}{\partial x_1 \partial x_{-1}} = -\sum_{m\in\mathbf{Z}_{\geq 0}} a_{n+1,m+1} e^{-\theta_m}, \quad n \in \mathbf{Z}_{\geq 0}, \tag{3.1}$$

where the matrix $A = (a_{n,m})$ is the transpose of the Cartan matrix for the infinite dimensional Lie algebra C_∞ [18].

The C_∞ 2D-Toda system (3.1) is transformed into the bilinear equations

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}} - 1\right)\tau_0 \cdot \tau_0 = \tau_1^2, \tag{3.2}$$

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}} - 1\right)\tau_n \cdot \tau_n = \tau_{n-1}\tau_{n+1}, \quad \text{for} \quad n \geq 1, \tag{3.3}$$

through the dependent variable transformation

$$\theta_0 = -\ln\frac{\tau_1^2}{\tau_0^2}, \qquad \text{and} \qquad \theta_n = -\ln\frac{\tau_{n+1}\tau_{n-1}}{\tau_n^2} \quad \text{for } n \geq 1. \tag{3.4}$$

Lemma 3.1. *The bilinear equations (3.2) and (3.3) have the N-soliton solution which is expressed as*

$$\tau_n = \det\left(\psi_{i,j}^{(n)}\right)_{1\leq i,j\leq 2N}, \tag{3.5}$$

where

$$\psi_{i,j}^{(n)} = c_{i,j} + (-1)^n \int_{-\infty}^{x_1} \varphi_i^{(n)}\varphi_j^{(-n)} dx_1, \tag{3.6}$$

$$\varphi_i^{(n)} = p_i^n e^{\xi_i}, \qquad \xi_i = p_i x_1 + \frac{1}{p_i}x_{-1} + p_i^3 x_3 + \frac{1}{p_i^3}x_{-3} + \cdots + \xi_{i0},$$

and $c_{i,j} = c_{j,i}$.

Proof. Imposing the C_∞ reduction $\tau_n = \tau_{-n}$, *i.e.*, folding the sequence of the τ-functions $\{\dots,\tau_{-2},\tau_{-1},\tau_0,\tau_1,\tau_2,\dots\}$ in τ_0, we have $\tau_{-1} = \tau_1$, $\tau_{-2} = \tau_2$, $\tau_{-3} = \tau_3$, ... [22, 19, 18, 23]. Thus we obtain the bilinear equations (3.2) and (3.3) from the 2D-Toda bilinear equation (2.3).

To impose the C_∞ reduction to the Gram-type determinant solution of the A_∞ 2D-Toda system, we impose the constraint $\hat{\varphi}_j^{(n)} = \varphi_j^{(n)}$, $c_{i,j} = c_{j,i}$, $M = 2N$ and $x_{2k} \equiv 0$ for every interger k. With this constraint, each element of the Gram-type determinant has the following property:

$$\begin{aligned}\psi_{i,j}^{(n)} &= c_{i,j} + (-1)^n \int_{-\infty}^{x_1} \varphi_i^{(n)} \varphi_j^{(-n)} dx_1 = c_{j,i} + (-1)^{-n} \int_{-\infty}^{x_1} \varphi_j^{(-n)} \varphi_i^{(n)} dx_1 \\ &= \psi_{j,i}^{(-n)} . \end{aligned} \tag{3.7}$$

Then the τ-function satisfies $\tau_n = \tau_{-n}$. Therefore the N-soliton solution of the C_∞ 2D-Toda system is expressed by the above Gram-type determinant. □

Lemma 3.2. *The bilinear equations*

$$-\left(\frac{1}{2} D_{x_1} D_{x_{-1}} - 1\right) \tau_0 \cdot \tau_0 = \tau_1^2 , \tag{3.8}$$

$$-\left(\frac{1}{2} D_{x_1} D_{x_{-1}} - 1\right) \tau_1 \cdot \tau_1 = \tau_1 \tau_0 , \tag{3.9}$$

have the N-soliton solution which is expressed as

$$\tau_n = \det \left(\psi_{i,j}^{(n)} \right)_{1 \le i,j \le 2N} , \tag{3.10}$$

where

$$\psi_{i,j}^{(n)} = c_{i,j} + (-1)^n \int_{-\infty}^{x_1} \varphi_i^{(n)} \varphi_j^{(-n)} dx_1 , \qquad \varphi_i^{(n)} = p_i^n e^{\xi_i} , \quad \xi_i = p_i x_1 + \frac{1}{p_i} x_{-1} + \xi_{i0} ,$$

and $c_{i,j} = \delta_{j,2N+1-i} \alpha_i$, $p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 0$.

Proof. Imposing a period 3-reduction $\tau_n = \tau_{n+3}$ to the sequence of the τ-functions $\{\ldots, \tau_3, \tau_2, \tau_1, \tau_0, \tau_1, \tau_2, \tau_3, \ldots\}$, it becomes $\{\ldots, \tau_0, \tau_1, \tau_1, \tau_0, \tau_1, \tau_1, \tau_0, \ldots\}$. Thus we obtain two bilinear equations (3.8) and (3.9).

In Lie algebraic terms, this corresponds to the reduction to the affine Lie algebra $A_2^{(2)}$ from the infinite dimensional Lie algebra C_∞.

To impose a period 3-reduction to the N-soliton solution, we add a constraint $p_i^3 + p_{2N+1-i}^3 = 0$, $(p_i \neq -p_{2N+1-i})$ for $i = 1, 2, \cdots, N$ and $c_{i,j} = \delta_{j,2N+1-i} \alpha_i$. To show $\tau_{n+3} = \tau_n$, we manipulate τ_n as follows:

$$\begin{aligned} \tau_n &= \det \left(\delta_{j,2N+1-i}\, \alpha_i + \frac{1}{p_i + p_j} \left(-\frac{p_i}{p_j} \right)^n e^{\xi_i + \xi_j} \right)_{1 \le i,j \le 2N} \\ &= e^{2 \sum_{k=1}^{2N} \xi_k} \det \left(\delta_{j,2N+1-i}\, \alpha_i \left(-\frac{p_j}{p_i} \right)^n e^{-\xi_i - \xi_j} + \frac{1}{p_i + p_j} \right)_{1 \le i,j \le 2N} \\ &= e^{2 \sum_{k=1}^{2N} \xi_k} \det \left(\delta_{j,2N+1-i}\, \alpha_i \left(-\frac{p_{2N+1-i}}{p_i} \right)^n e^{-\xi_i - \xi_{2N+1-i}} + \frac{1}{p_i + p_j} \right)_{1 \le i,j \le 2N} \end{aligned}$$

Since $\left(-\frac{p_{2N+1-i}}{p_i} \right)^3 = 1$, $\tau_{n+3} = \tau_n$ is satisfied. □

Lemma 3.3. *The τ-function τ_1 of the bilinear equations (3.8) and (3.9) is written in the form of pfaffian*

$$\tau_1 = \frac{1}{2^{2N} \prod_{k=1}^{2N} p_k} \tau^2 , \qquad \tau = \mathrm{pf}(1, 2, ..., 2N-1, 2N) , \tag{3.11}$$

where $\mathrm{pf}(i,j) = c_{i,j} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}$, $\xi_i = p_i^{-1}x_{-1} + p_i x_1 + \xi_i^0$, $c_{i,j} = \delta_{j,2N+1-i}c_i$, $c_i = -c_{2N+1-i}$, $p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 0$.

Proof. Let $\alpha_i = \alpha_{2N+1-i}$. For $n = 1$,

$$\begin{aligned}
\tau_1 &= \det\left(\psi_{i,j}^{(1)}\right)_{1\le i,j\le 2N} = \det\left(\delta_{j,2N+1-i}\,\alpha_i - \frac{1}{p_i+p_j}\frac{p_i}{p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \frac{1}{2^{2N}\prod_{k=1}^{2N}p_k}\det\left(2\delta_{j,2N+1-i}\,\alpha_i\frac{p_j^2}{p_i} - \frac{2p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \frac{1}{2^{2N}\prod_{k=1}^{2N}p_k}\det\left(2\delta_{j,2N+1-i}\,\alpha_i\frac{p_{2N+1-i}^2}{p_i} - \frac{2p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \frac{1}{2^{2N}\prod_{k=1}^{2N}p_k}\det\left(c_{i,j} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j} - e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N},
\end{aligned}$$

where $c_{i,j} = 2\delta_{j,2N+1-i}\,\alpha_i\frac{p_{2N+1-i}^2}{p_i}$. Then we note

$$c_{i,j} = 2\delta_{j,2N+1-i}\,\alpha_i\,\frac{p_{2N+1-i}^2}{p_i} = -2\delta_{i,2N+1-j}\,\alpha_{2N+1-i}\,\frac{p_i^2}{p_{2N+1-i}} = -c_{j,i}\,.$$

Introducing $c_i = 2\alpha_i\frac{p_{2N+1-i}^2}{p_i}$, we can write as $c_{i,j} = \delta_{j,2N+1-i}c_i$ and $c_i = -c_{2N+1-i}$.

Since the $2N\times 2N$ matrix $\left(c_{i,j} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}$ is skew-symmetric, we obtain

$$\tau_1 = \frac{1}{2^{2N}\prod_{k=1}^{2N}p_k}\left[\mathrm{pf}(1,2,\ldots,2N-1,2N)\right]^2,$$

where $\mathrm{pf}(i,j) = c_{i,j} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}$. Here we used the formula [21]

$$\det(a_{i,j} - y_i y_j)_{1\le i,j\le 2N} = \det(a_{i,j})_{1\le i,j\le 2N} = \left[\mathrm{pf}(1,2,\ldots,2N)\right]^2,$$

where $a_{i,j} = -a_{j,i}$, $a_{i,i} = 0$, $\mathrm{pf}(i,j) = a_{i,j}$. $\square$

Letting $f = \tau_1$ and $g = \tau_0$, we obtain the bilinear equations (2.20) and (2.21) from the bilinear equations (3.9) and (3.8). As is shown in [18], the period 3-reduction of C_∞ 2D-Toda system gives the same result as the period 3-reduction of B_∞ 2D-Toda after the relabelling indexes.

References

[1] Ostrovsky L A 1978 Nonlinear internal waves in a rotating ocean, *Oceanology* **18**, 119–125

[2] Stepanyants Y A 2006 On stationary solutions of the reduced Ostrovsky equation: Periodic waves, compactons and compound solitons, *Chaos, Solitons & Fractals* **28**, 193–204

[3] Hunter J 1990 Numerical solutions of some nonlinear dispersive wave equations, *Lectures in Appl. Math.* **26**, 301–316

[4] Vakhnenko V A 1992 Solitons in a nonlinear model medium, *J. Phys. A: Math. Gen.* **25**, 4181–4187

[5] Vakhnenko V O and Parkes E J 1998 The two loop soliton solution of the Vakhnenko equation, *Nonlinearity* **11**, 1457–1464

[6] Morrison A J, Vakhnenko V O and Parkes E J 1999 The N loop soliton solution of the Vakhnenko equation, *Nonlinearity,* **12**, 1427–1437

[7] Vakhnenko V O and Parkes E J 2002 The calculation of multi-soliton solutions of the Vakhnenko equation by the inverse scattering method, *Chaos, Solitons & Fractals* **13**, 1819–1826

[8] Liu Y, Pelinovsky D and Sakovich A 2010 Wave Breaking in the Ostrovsky-Hunter Equation, *SIAM J. Math. Anal.* **42**, 1967–1985

[9] Hone A N W and Wang J P 2003 Prolongation algebras and Hamiltonian operators for peakon equations, *Inverse Problems* **19**, 129–145

[10] Matsuno Y 2006 Cusp and loop soliton solutions of short-wave models for the Camassa-Holm and Degasperis-Procesi equations, *Phys. Lett. A* **359**, 451–457

[11] Degasperis A and Procesi M 1999 Asymptotic integrability, in *Symmetry and Perturbation Theory*, edited by A. Degasperis and G. Gaeta, World Scientific 23–37

[12] Matsuno Y 2005 Multisoliton solution of the Degasperis-Procesi equation and their peakon limit, *Inverse Problems* **21**, 1553–1570

[13] Matsuno Y 2005 The N-soliton solution of the Degasperis-Procesi equation, *Inverse Problems* **21**, 2085–2101

[14] Leznov A N and Saveliev M V 1979 Representation of zero curvature for the system of nonlinear partial differential equations $x_{a,z,\bar{z}} = \exp(kx)_a$ and its integrability, *Lett. Math. Phys.* **3**, 489–494

[15] Mikhailov A V 1979 On the integrability of a two-dimensional generalization the Toda lattice, *JETP Lett.* **30**, 414–418

[16] Mikhailov A V 1981 The reduction problem and the inverse scattering method, *Physica* **3D**, 73–117

[17] Mikhailov A V 1981 Two-dimensional generalized Toda lattice, *Commun. Math. Phys.* **79**, 473–488

[18] Nimmo J J C and Willox R 1997 Darboux transformations for the two-dimensional Toda system, *Proc. R. Soc. Lond.* **453**, 2497–2525

[19] Ueno K and Takasaki K 1984 Toda lattice hierarchy, *Adv. Stud. Pure Math.* **4**, 1–95

[20] Babich V M, Matveev V B and Sall M A 1986 Binary Darboux transformation for the Toda lattice, *J. Math. Sci.* **35**, 2582–2589

[21] Hirota R 2004 *The Direct Method in Soliton Theory*, Cambridge University Press.

[22] Jimbo M and Miwa T 1983 Solitons and infinite dimensional Lie algebras, *Publ. RIMS. Kyoto Univ.* **19**, 943–1001

[23] Willox R 2005 On a generalized Tzitzeica equation, *Glasgow Math. J.* **47A**, 221–231

[24] Nimmo J J C and Ruijsenaars S N M 2009 Tzitzeica solitons versus relativistic Calogero-Moser three-body clusters, *J. Math. Phys.* **50**, 043511 (31pp)

[25] Tzitzeica G 1910 Sur une nouvelle classe de surfaces, *C. R. Acad. Sci.* **150**, 955–956

[26] Tzitzeica G 1910 Sur une nouvelle classe de surfaces, *C. R. Acad. Sci.* **150**, 1227–1229

[27] Dodd R K and Bullough R K 1977 Polynomial conserved densities for the sine-Gordon equations, *Proc. R. Soc. London, Ser. A*, **352**, 481–503

[28] Date E, Jimbo M, Kashiwara M and Miwa T 1982 Transformation groups for soliton equations IV. A new hierarchy of soliton equations of KP-type, *Physica* **4D**, 343–365

[29] Date E, Jimbo M, Kashiwara M and Miwa T 1981 KP hierarchies of orthogonal and symplectic type –Transformation groups for soliton equations VI–, *J. Phys. Soc. Japan* **50**, 3813–3818

[30] Date E, Jimbo M, Kashiwara M and Miwa T 1982 Transformation groups for soliton equations– Euclidean Lie algebras and reduction of the KP hierarchy, *Publ. RIMS. Kyoto Univ.* **18**, 1077–1110

[31] Hirota R 1989 Soliton Solutions to the BKP Equations. I. The Pfaffian technique, *J. Phys. Soc. Japan* **58**, 2285–2296

[32] Hirota R 1989 Soliton Solutions to the BKP Equations. II. The Integral Equation, *J. Phys. Soc. Japan* **58**, 2705–2712

[33] Ito M 1980 An Extension of Nonlinear Evolution Equations of the K-dV (mK-dV) Type to Higher Orders, *J. Phys. Soc. Japan* **49**, 771–778

[34] Lambert F, Loris I, Springael J and Willox R 2001 On the Hirota Representation of Soliton Equations with One Tau-Function, *J. Phys. Soc. Japan* **70**, 605–608

[35] Feng B-F, Maruno K and Ohta Y 2010 Integrable discretizations for the short-wave model of the Camassa-Holm equation, *J. Phys. A:* **43**, 265202 (13pp)